

Table 3 Distribution zone, utility, 6-month body weight and dressing percentage of breeds/populations investigated

Utility	Breed	Distribution		6-month Weight (kg)	Dressing (%)	Ref.
		Region	State			
Meat	Bandur/Mandya	SP	Karnataka	17.28	54.5	Singh et al. 2005
	Deccani	SP	Maharashtra, Andhra Pradesh,	19.26	52.65	Singh et al. 2005
			Karnataka			
	Garole	E	West Bengal	8.97	55.87	Singh et al. 2005 Banerjee et al. 2010
	Madgyal	SP	Maharashtra	25-30	NA	Anonymous
Nellore	SP	Andhra Pradesh	17.81	58.22	Singh et al. 2005	
Carpet Wool	Chokla	NWASA	Rajasthan	13.46	NA	Acharya 1982
	Magra	NWASA	Rajasthan	20.14	44.75	Acharya 1982
	Nali	NWASA	Rajasthan	13.30	48.25	Acharya 1982
Meat, Carpet Wool	Ganjam	E	Orissa	12.92	NA	Pattanayak et al. 2003
	Malpura	NWASA	Rajasthan	24.58	59.13	Singh et al. 2005
	Muzaffarnagri	NWASA	Uttar Pradesh, Uttarakhand	21.17	57.26	Singh et al. 2005

Note: SP-Southern Peninsular; NWASA-Northwestern Arid and Semi Arid; E-Eastern; NA-Not Available